

PS

Weed



Net Wt. 20 Lbs. (9.07 kg) **KEEP OUT OF REACH OF CHILDREN** CAUTION **Statement of Practical Treatment IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists. IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting. See back panel for Precautionary Statements.





1 Control -raniles TO TO T ATTITUTO FOR LAWNS

CONTROLS DANDELIONS, PLANTAINS, CLOVER, CHICKWEED **AND OVER 200 OTHER WEEDS** Treats 10,000 Sq. Ft.

ACTIVE INGREDIENTS: Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid... ** Dimethýlamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid......
*** Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)..... 0.794% 0.183% Other Ingredients.... Total.... 100.000% THIS PRODUCT CONTAINS: * 27.36 lb.s 2,4-Dichlorophenoxyacetic acid equivalent per ton or 1.368% * 13.12 lbs. 2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.656% *** 3.04 lbs. 3,6-dichloro-o-anisic acid equivalent per ton or 0.152% Isomer Specific by AOAC Methods EPA Reg. No. 2217-657-9198 EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A Trimec® is a registered trademark of PBI/GORDON CORPORATION; (Underlined letter used is first letter of run code on end of bag Fortify is a registered trademark of The Andersons Agriservices, Inc.

Ί

long pants, socks and shoes.

ENVIRONMENTAL HAZARDS This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target area. Do not contaminate domestic or irrigation waters.

not enter treatment areas until spray has dried or dust has settled.

production, or for research purposes. _____ STORAGE: locked storag feedstuffs. discard in tro

Do not apply this product to St. Augustinegrass, carpetgrass, dichondra, ornamentals, or lawns containing desirable clovers. Avoid sprading granules onto landscape, plantings, flowers, vegetables or ornamentals. Do not spread over exposed roots of trees and ornamentals. Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species. Treated areas may be seeded 3 to 4 weeks after application. Application of this product to newly seeded areas should be delayed until the third mowing.

Aste

Austrian fieldcress Bedstraw Beggarticks Betony, Florida Bindweed, field Bird vetch Bitter wintercress Bittercress, hairy Bitterweed Black-eyed Susan Black medic Black mustard Blackseed plantain Blessed thistle Blue lettuce Blue vervain Bracted plantain Brassbuttons Bristly oxtongue Broadleaf dock Broadleaf plantain Buckhorn Buckhorn plantain Bulbous buttercop Bull thistle Burdock Burweed Buttercup Buttonweed Carolina geranium Carpetweed Chickweed, mouseear Chicory Cinquefoil Clover, crimson Clover, hop Clover, red Clover, strawberry Clover, sweet Clover, white Cockle Cocklebur Common mullein Creeping jenny

Weed Control Granules

Weed Control Førtify® Granules

STOP! Read the entire label first. Observe all precautions and follow directions carefully.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS **CAUTION:** Harmful if swallowed. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt,

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

DIRECTIONS FOR USE It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed

| · · · · · · · · · · · · · · · · · · · | | | |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|--|--|
| : Store in original container and keep any bags containing unused product in a age area inaccessible to children and domestic animals. Do not store near food or PRODUCT DISPOSAL: Securely wrap bag in several layers of newspaper and rash. CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash. | | | |
| Tash. CONTAINER DISPOSAL. Do noi reuse bag. Discura bag in Irash. | | | |
| USE PRECAUTIONS | | | |

WHERE TO USE

Fortify Weed Control is a TRIMEC® complex, a combination of three proven weed killers - 2,4-D, Mecoprop and Dicamba. Together, they control a wide range of lawn weeds including dandelion, chickweed, knotweed, plantains, henbit, spurge and many others found in home lawns. Established turf of Kentucky bluegrass, bentgrass, bahiagrass, centipedegrass, fescues, bermudagrass, zoysiagrass, and perennial ryegrass may be treated.

WEEDS CONTROLLED

Dandelion Dogfennel Dollarweed False dandelior False flax Florida pusley Frenchweed Goathead Ground Ivy Hawkweed Healall Heathaster Hedge mustard Hemp Henbit Hoary cress Hoary plantain Hoary vervain Horsenettle Knotweed Lambsquarters Lespedeza Mallow Matchweed Moringglory Musk thistle Mustard Narrowleaf plantain Narrowleaf vetch Oxalis Parsnip Pennycress Catsear Chickweed, common Pennywort Peppergrass Pigweed Plantain Poison Ivy Poison oal Pokeweed Prairie sunflower Prickly lettuce Prickly sida Prostrate knotweed Prostrate pigweed Prostrate spurge Prostrate vervain

Cudweed Curly dock Daisy, oxeye Puncturevine

Ragweed

Red sorrel

Purslane, common


Redroot pigweed Rough cinquefoil Russian pigweed Russian thistle Scotch thistle Shepherspurse Slender plantain Smooth pigweed Sorrel Sowthistle Speedwell Spiny cocklebur Spiny sowthistle Spotted spurge Spurweed Stinging nettle Tall nettle Tansy ragwort Tansy mustard Thistle Tumble mustard Tumble pigweed Velvetleaf Venice mallow Virginia buttonweed Virginia creeper Water pennywort White mustard Wild aster Wild carrot Wild garlic Wild geranium Wild lettuce Wild marigold Wild mustard Wild onion Wild parsnip Wild radish Wild strawberry Wild vetch Woodsorrel Woolly croton Woolly morningglory Woolly plantain Yarrow Yellow rocket and other broadleaf weeds

SPREADING SUGGESTIONS

It is important to follow the directions as indicated on this quality lawn care product as over-application may cause injury to your lawn. Measure the size of area to be treated and do not over or under apply. Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Calibrate spreader prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing uniformly. Close spreader when filling, stopping, or turning. Do not apply by hand or with a hand-held spreader.

BROADCAST SPREADERS: For best results, apply in a circular path starting on the outside and working toward the center (see Path A below). A little overlap of product will ensure complete coverage.

DROP SPREADERS: First apply 2 border strips along the edge of each end of the lawn (see Path B below). Then apply product along the length of the lawn, overlapping slightly and shutting off the spreader before turning as you reach the edge of the border strips.



SUGGESTED SPREADER SETTINGS 2.0 Lbs. of Product/1000 Sq. Ft.

BROADCAST SPREADERS:

| Earthway 2000A Precision SB50 Republic EZ540/75450 Scotts SpeedyGreen Scotts EasyGreen | 6 2 1/2 4 |
|----------------------------------------------------------------------------------------------------|-----------------|
| Earthway 7350 | |

| | - |
|----------------------|---|
| Precision FH22 | 7 |
| Republic EZ530/75350 | 8 |
| Scotts PF4 | |
| Scotts AccuGreen | 5 |
| | |

The above spreader settings are approximate. Calibrate your spreader on 100 square feet and adjust at the rate of 1 pound per 500 square feet.

WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NOT OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is the Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

Distributed by: **Free Flow Fertilizer** P. O. Box 119 Maumee, OH 43537

FRTBK2

©2000 T.A.

